

2. Introduction to Continued Fractions (16 July 2019).

2.1. What is Euclidean algorithm and how it is related to continued fractions of rational numbers?

2.2. Formal infinite continued fraction.

$$[a_0; a_1, a_2, \dots, a_\nu, \dots], \quad a_0 \in \mathbb{Z}, \quad a_j \in \mathbb{Z}_+, j = 1, 2, 3, \dots \quad (1)$$

a_j - partial quotients,

$$\frac{p_\nu}{q_\nu} = [a_0; a_1, a_2, \dots, a_\nu] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots + \frac{1}{a_\nu}}}, \quad (p_\nu, q_\nu) = 1 - \text{convergents.}$$

2.3. Recursive formulas for the convergents' numerators and denominators.

$$p_{\nu+1} = a_{\nu+1}p_\nu + p_{\nu-1}, \quad q_{\nu+1} = a_{\nu+1}q_\nu + q_{\nu-1}, \quad p_\nu q_{\nu-1} - q_\nu p_{\nu-1} = (-1)^{\nu-1}.$$

2.4. The value of continued fraction (1). Prove that

- a. $\frac{p_{2\nu}}{q_{2\nu}}$ is an increasing sequence;
- b. $\frac{p_{2\mu+1}}{q_{2\mu+1}}$ is a decreasing sequence;
- c. $\frac{p_{2\nu}}{q_{2\nu}} < \frac{p_{2\mu+1}}{q_{2\mu+1}}$ for all μ, ν ;
- d. $\left| \frac{p_\nu}{q_\nu} - \frac{p_{\nu+1}}{q_{\nu+1}} \right| = \frac{1}{q_\nu q_{\nu+1}}$;
- e. there exists $\lim_{\nu \rightarrow \infty} \frac{p_\nu}{q_\nu}$ which is called the value of continued fraction (1).

2.5. For every real number α there exists a continued fraction of the form (1) which value is α .

2.6. Problem of uniqueness. Prove that every irrational number has the unique representation as a value of a continued fraction of the form (1). What happens with rational numbers, and what is the correct statement about uniqueness for rationals?

2.7. Prove that

$$\|q_\nu \alpha\| = \frac{1}{q_\nu(\alpha_{\nu+1} + \alpha_\nu^*)},$$

where

$$\alpha_{\nu+1} = [a_{\nu+1}; a_{\nu+2}, a_{\nu+3}, \dots], \quad \alpha_\nu^* = [0; a_\nu, a_{\nu-1}, \dots, a_1].$$

2.8. **Lagrange Theorem.** α is a quadratic irrationality if and only if its continued fraction is eventually periodic.

2.9 **Zaremba's Conjecture.**

$$\forall q \in \mathbb{Z}_+ \quad \exists a : (a, q) = 1 \text{ such that in its c.f. expansion } a_j \leq 5, \forall j.$$

(We will not prove it.)

Exercises.

1. Prove that for any α and for any ν one has $q_\nu \geq \left(\frac{1+\sqrt{5}}{2}\right)^{\nu-1}$.

2. **Valen's theorem.** For any ν either

$$q_\nu \|q_\nu \xi\| < 1/2,$$

or

$$q_{\nu+1} \|q_{\nu+1} \xi\| < 1/2$$

holds.

3. Suppose that in (1) $a_0 \geq 1$. Prove that $\frac{p_n}{p_{n-1}} = [a_n; a_{n-1}, \dots, a_0]$.

4. Prove that

a. $\sqrt{d^2 + 1} = [d; \overline{2d}]$;

b. $\sqrt{d^2 + 2} = [d; \overline{d, 2d}]$;

c. $\underbrace{[2; 2, \dots, 2]}_n = \frac{(1+\sqrt{2})^{n+1} - (1-\sqrt{2})^{n+1}}{(1+\sqrt{2})^n - (1-\sqrt{2})^n}$

5. Prove that each rational number $\frac{a}{b}$ can be represented in a form

$$b_0 - \frac{1}{b_1 - \frac{1}{b_2 - \dots - \frac{1}{b_\nu}}} \tag{2}$$

with $b_j \geq 2, j = 1, 2, \dots, \nu$.

6. Prove Zaremba's Conjecture for

a. $q = F_n$ - Fibonacci numbers;

b. $q = 2^n$;

c. for representation of rationals as continues fractions (2), that is, you should prove that for any $q \in \mathbb{Z}_+$ there exists $a \in \mathbb{Z}$ such that $(a, q) = 1$ and in the decomposition

$$b_0 - \frac{1}{b_1 - \frac{1}{b_2 - \dots - \frac{1}{b_\nu}}}$$

we have $b_j \leq 5 \forall j$.